

Complexes of Nonseparating Curves and Mapping Class Groups

Elmas Irmak

Abstract

Let R be a compact, connected, orientable surface of genus g , Mod_R^* be the extended mapping class group of R , $\mathcal{C}(R)$ be the complex of curves on R , and $\mathcal{N}(R)$ be the complex of nonseparating curves on R . We prove that if $g \geq 2$ and R has at most $g - 1$ boundary components, then a simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ is superinjective if and only if it is induced by a homeomorphism of R . We prove that if $g \geq 2$ and R is not a closed surface of genus two then $Aut(\mathcal{N}(R)) = Mod_R^*$, and if R is a closed surface of genus two then $Aut(\mathcal{N}(R)) = Mod_R^*/\mathcal{C}(Mod_R^*)$. We also prove that if $g = 2$ and R has at most one boundary component, then a simplicial map $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is superinjective if and only if it is induced by a homeomorphism of R . As a corollary we prove some new results about injective homomorphisms from finite index subgroups to Mod_R^* . The last two results complete the author's previous results to connected orientable surfaces of genus at least two.

MSC: 57M99, 20F38.

Keywords: Mapping class groups; Surfaces; Complex of curves.

1 Introduction

Let R be a compact, connected, orientable surface of genus g with p boundary components. The extended mapping class group, Mod_R^* , of R is the group of isotopy classes of all (including orientation reversing) homeomorphisms of R . Let $\mathcal{A}$ denote the set of isotopy classes of nontrivial simple closed curves on R . The *complex of curves*, $\mathcal{C}(R)$, on R is an abstract simplicial complex, introduced by Harvey [2], with vertex set $\mathcal{A}$ such that a set of n vertices $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ forms an $n - 1$ simplex if and only if $\alpha_1, \alpha_2, \dots, \alpha_n$ have pairwise disjoint representatives. Let $\mathcal{B}$ denote the set of isotopy classes of nonseparating simple closed curves on R . The *complex of nonseparating curves*, $\mathcal{N}(R)$, is the subcomplex of $\mathcal{C}(R)$ with the vertex set $\mathcal{B}$ such that a set of n vertices $\{\beta_1, \beta_2, \dots, \beta_n\}$ forms an $n - 1$ simplex if and only if $\beta_1, \beta_2, \dots, \beta_n$ have pairwise disjoint representatives. The main results of the paper are the following:

Theorem 1.1 *Suppose that $g \geq 2$ and R has at most $g - 1$ boundary components. Then a simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ is superinjective if and only if λ is induced by a homeomorphism of R .*

Theorem 1.2 *Suppose that $g \geq 2$. If R is not a closed surface of genus two, then $Aut(\mathcal{N}(R)) = Mod_R^*$. If R is a closed surface of genus 2, then $Aut(\mathcal{N}(R)) = Mod_R^*/\mathcal{C}(Mod_R^*)$.*

Supported by a Rackham Faculty Fellowship, The Rackham Graduate School, University of Michigan.

Theorem 1.3 Suppose $g = 2$ and $p \leq 1$. A simplicial map $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is superinjective if and only if λ is induced by a homeomorphism of R .

Theorem 1.4 Let K be a finite index subgroup of Mod_R^* and f be an injective homomorphism $f : K \rightarrow \text{Mod}_R^*$. If $g = 2$ and $p = 1$ then f has the form $k \rightarrow hkh^{-1}$ for some $h \in \text{Mod}_R^*$ and f has a unique extension to an automorphism of Mod_R^* . If R is a closed surface of genus 2, then f has the form $k \rightarrow hkh^{-1}i^{m(k)}$ for some $h \in \text{Mod}_R^*$ where m is a homomorphism $K \rightarrow \mathbb{Z}_2$ and i is the hyperelliptic involution on R .

In section 2, we give some properties of the superinjective simplicial maps of $\mathcal{N}(R)$. In section 3, we give some properties of the superinjective simplicial maps of $\mathcal{C}(R)$ and we prove Theorem 1.3 and Theorem 1.4. In section 4, we prove that a superinjective simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ extends to a superinjective simplicial map on $\mathcal{C}(R)$, and we prove Theorem 1.1 and Theorem 1.2.

Theorem 1.3 and Theorem 1.4 complete the author's previous results given in [3] and [4] to connected orientable surfaces of genus at least two. These theorems were motivated by the work of Ivanov [5], and Ivanov and McCarthy [7]. For automorphism groups of some complexes related to complex of nonseparating curves $\mathcal{N}(R)$, see [11], [13].

2 Properties of Superinjective Simplicial Maps of $\mathcal{N}(R)$

A *circle* on R is a properly embedded image of an embedding $S^1 \rightarrow R$. A circle on R is said to be *nontrivial* (or *essential*) if it doesn't bound a disk and it is not homotopic to a boundary component of R . Let C be a collection of pairwise disjoint circles on R . The surface obtained from R by cutting along C is denoted by R_C . A nontrivial circle a on R is called *nonseparating* if the surface R_a is connected. Let α and β be two vertices in $\mathcal{N}(R)$. The *geometric intersection number* $i(\alpha, \beta)$ is defined to be the minimum number of points of $a \cap b$ where $a \in \alpha$ and $b \in \beta$. A simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ is called **superinjective** if the following condition holds: if α, β are two vertices in $\mathcal{N}(R)$ such that $i(\alpha, \beta) \neq 0$, then $i(\lambda(\alpha), \lambda(\beta)) \neq 0$.

Lemma 2.1 Suppose $g \geq 2$ and $p \geq 0$. A superinjective simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ is injective.

Proof. Let α and β be two distinct vertices in $\mathcal{N}(R)$. If $i(\alpha, \beta) \neq 0$, then $i(\lambda(\alpha), \lambda(\beta)) \neq 0$, since λ preserves nondisjointness. So, $\lambda(\alpha) \neq \lambda(\beta)$. If $i(\alpha, \beta) = 0$, then, since $g \geq 2$ and $p \geq 0$, we can choose a vertex γ of $\mathcal{N}(R)$ such that $i(\gamma, \alpha) = 0$ and $i(\gamma, \beta) \neq 0$. Then $i(\lambda(\gamma), \lambda(\alpha)) = 0$ and $i(\lambda(\gamma), \lambda(\beta)) \neq 0$. So, $\lambda(\alpha) \neq \lambda(\beta)$. Hence λ is injective. $\square$

Let P be a set of pairwise disjoint circles on R . P is called a *pair of pants decomposition* of R , if R_P is a disjoint union of genus zero surfaces with three boundary components, pairs of pants. A pair of pants of a pants decomposition is the image of one of these connected components under the quotient map $q : R_P \rightarrow R$. The image of the boundary of this component is called the *boundary of the pair of pants*. A pair of pants is called *embedded* if the restriction of q to the corresponding component of R_P is an embedding. An ordered set $(a_1, \dots, a_{3g-3+p})$ is called an *ordered pair of pants decomposition* of R if $\{a_1, \dots, a_{3g-3+p}\}$ is a pair of pants decomposition of R .

Lemma 2.2 Suppose $g \geq 2$ and $p \geq 0$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Let P be a pair of pants decomposition consisting of nonseparating circles on R . Then λ maps the set of isotopy classes of elements of P to the set of isotopy classes of elements of a pair of pants decomposition P' of R .

Proof. The set of isotopy classes of elements of P forms a top dimensional simplex, Δ , in $\mathcal{N}(R)$. Since λ is injective, it maps Δ to a top dimensional simplex Δ' in $\mathcal{N}(R)$. A set of pairwise disjoint representatives of the vertices of Δ' is a pair of pants decomposition P' of R . $\square$

Let P be a pair of pants decomposition of R . Let a and b be two distinct elements in P . Then a is called *adjacent* to b w.r.t. P iff there exists a pair of pants in P which has a and b on its boundary.

Remark: Let P be a pair of pants decomposition of R . Let $[P]$ be the set of isotopy classes of elements of P . Let $\alpha, \beta \in [P]$. We say that α is adjacent to β w.r.t. $[P]$ if the representatives of α and β in P are adjacent w.r.t. P . By Lemma 2.2, λ gives a correspondence on the isotopy classes of elements of pair of pants decompositions consisting of nonseparating circles on R . $\lambda([P])$ is the set of isotopy classes of elements of a pair of pants decomposition which corresponds to P , under this correspondence.

Lemma 2.3 *Suppose $g \geq 2$ and $p \geq 0$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Let P be a pair of pants decomposition consisting of nonseparating circles on R . Then λ preserves the adjacency relation for two circles in P , i.e. if $a, b \in P$, a is adjacent to b w.r.t. P and $[a] = \alpha, [b] = \beta$, then $\lambda(\alpha)$ is adjacent to $\lambda(\beta)$ w.r.t. $\lambda([P])$.*

Proof. Let P be a pair of pants decomposition consisting of nonseparating circles on R . If $g = 2$ and $p \leq 1$, then every element in $\lambda([P])$ is adjacent to any other element in $\lambda([P])$, so the lemma is clear. For the other cases; Let a, b be two adjacent circles in P and $[a] = \alpha, [b] = \beta$. By Lemma 2.2, we can choose a pair of pants decomposition, P' , such that $\lambda([P]) = [P']$. Let P_o be a pair of pants of P , having a and b on its boundary. P_o is an embedded pair of pants. There are two possible cases for P_o , depending on whether a and b are the boundary components of another pair of pants or not. For each of these cases, we show how to choose a circle c which essentially intersects a and b and does not intersect any other circle in P in Figure 1.




Figure 1: Two possible cases for P_o

Let $\gamma = [c]$. Assume that $\lambda(\alpha)$ and $\lambda(\beta)$ do not have adjacent representatives. Since $i(\gamma, \alpha) \neq 0$ and $i(\gamma, \beta) \neq 0$, we have $i(\lambda(\gamma), \lambda(\alpha)) \neq 0$ and $i(\lambda(\gamma), \lambda(\beta)) \neq 0$ by superinjectivity. Since $i(\gamma, [e]) = 0$ for all e in $P \setminus \{a, b\}$, we have $i(\lambda(\gamma), \lambda([e])) = 0$ for all e in $P \setminus \{a, b\}$. But this is not possible because $\lambda(\gamma)$ has to intersect geometrically essentially with some isotopy class other than $\lambda(\alpha)$ and $\lambda(\beta)$ in the image pair of pants decomposition to be able to make essential intersections with $\lambda(\alpha)$ and $\lambda(\beta)$. This is a contradiction to the assumption that $\lambda(\alpha)$ and $\lambda(\beta)$ do not have adjacent representatives. $\square$




Figure 2: Curve configurations

Let P be a pair of pants decomposition of R . A curve $x \in P$ is called a 4 -curve in P if there exist four distinct circles in P , which are adjacent to x w.r.t. P . Note that in a pants decomposition every curve is adjacent to at most 4 curves.

Lemma 2.4 *Suppose $g \geq 2$ and $p \geq 0$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map and α, β, γ be distinct vertices in $\mathcal{N}(R)$ having pairwise disjoint representatives which bound a pair of pants in R . Then $\lambda(\alpha), \lambda(\beta), \lambda(\gamma)$ are distinct vertices in $\mathcal{N}(R)$ having pairwise disjoint representatives which bound a pair of pants in R .*

Proof. Let a, b, c be pairwise disjoint representatives of α, β, γ respectively. If R is a closed surface of genus 2 then the statement is obvious. Consider the case when $g = 2, p = 1$. We complete $\{a, b, c\}$ to a pair of pants decomposition $P = \{a, b, c, a_1\}$ consisting of nonseparating circles as shown in Figure 2 (i). Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c', a'_1 be the representatives of $\lambda([a]), \lambda([b]), \lambda([c]), \lambda([a_1])$ in P' respectively. Since a is adjacent to b w.r.t. P , a' is adjacent to b' w.r.t. P' . Then there is a pair of pants Q_1 in P' having a' and b' on its boundary. Let x be the other boundary component of Q_1 . If $x = c'$ then we are done. Assume that $x \neq c'$. Then x can't be a' , since otherwise b' would be a separating circle which is a contradiction. Similarly, x can't be b' . Then x is either a'_1 or it is the boundary component of R .

Assume x is a boundary component of R . Then, since a is adjacent to c, a_1 w.r.t. P , a' is adjacent to c', a'_1 w.r.t. P' and so, there is a pair of pants Q_2 in P' having a' and c', a'_1 on its boundary. Similarly, since b is adjacent to c, a_1 w.r.t. P , b' is adjacent to c', a'_1 w.r.t. P' and so, there is a pair of pants Q_3 in P' having b' and c', a'_1 on its boundary. Then it is clear that $R = Q_1 \cup Q_2 \cup Q_3$. Now we consider the circles a_2, a_3 which are as shown in Figure 2 (i). Since a_2 intersects b essentially and a_2 is disjoint from a , we can choose a representative a'_2 of $\lambda([a_2])$ such that there exists an essential arc w of a'_2 in Q_1 which starts and ends on b' and which does not intersect $a' \cup \partial R$. Since a_3 is disjoint from b and a_2 , there exists a representative a'_3 of $\lambda([a_3])$ such that a'_3 is disjoint from $b' \cup w$. But then a'_3 could be isotoped so that it is disjoint from a' , since a' is a boundary component of a regular neighborhood of $b' \cup w$ in Q_1 . This is a contradiction since $i([a], [a_3]) \neq 0$ and so $i(\lambda([a]), \lambda([a_3])) \neq 0$. So, x can not be a boundary component of R .

Assume that $x = a'_1$. Then, since a is adjacent to c w.r.t. P , a' is adjacent to c' w.r.t. P' and so, there is a pair of pants Q_2 in P' having a' and c' on its boundary. Let y be the other boundary component of Q_2 . If $y = c'$, then a' would be a separating circle, which is a contradiction. If y is the boundary component of R , then since c is adjacent to b, a_1 w.r.t. P , c' is adjacent to b', a'_1 in P' and so, there is a pair of pants Q_3 in P' having b', c', a'_1 on its

boundary. Then it is clear that $R = Q_1 \cup Q_2 \cup Q_3$. Now we consider the circles a_2, a_4 which are as shown in Figure 2 (i). Since a_4 intersects a essentially and a_4 is disjoint from c , we can choose a representative a'_4 of $\lambda([a_4])$ such that there exists an essential arc w of a'_4 in Q_2 which starts and ends on a' and which does not intersect $c' \cup \partial R$. Now, we consider a_2 ; a_2 is disjoint from a and a_4 . Then there exists a representative a'_2 of $\lambda([a_2])$ such that a'_2 is disjoint from $a' \cup w$. But then a'_2 could be isotoped so that it is disjoint from c' , since c' is a boundary component of a regular neighborhood of $a' \cup w$ in Q_2 . This is a contradiction since $i([c], [a_2]) \neq 0$ and so $i(\lambda([c]), \lambda([a_2])) \neq 0$. So, y can not be the boundary component of R . If $y = a'_1$, then there exists a pair of pants Q_3 having b', c' and the boundary component of R as its boundary components. Then we have $R = Q_1 \cup Q_2 \cup Q_3$. Now, we consider the circles a_3, a_4 . Since a_3 intersects c essentially and a_3 is disjoint from b , we can choose a representative a'_3 of $\lambda([a_3])$ such that there exists an essential arc w of a'_3 in Q_3 which starts and ends on c' and which does not intersect $b' \cup \partial R$. Now, we consider a_4 ; a_4 is disjoint from c and a_3 . Then there exists a representative a'_4 of $\lambda([a_4])$ such that a'_4 is disjoint from $c' \cup w$. But then a'_4 could be isotoped so that it is disjoint from b' , since b' is a boundary component of a regular neighborhood of $c' \cup w$ in Q_3 . This is contradiction since $i([a_4], [b]) \neq 0$ and so $i(\lambda([a_4]), \lambda([b])) \neq 0$. Hence, we conclude that $y = b'$. Then a', b', c' bound a pair of pants which completes the proof for $g = 2, p = 1$.

For the remaining cases we take a subsurface of genus two with two boundary components, N , of R , containing nonisotopic circles a, b, c, a_1, a_2 as shown in Figure 2 (ii). Then we complete $\{a, b, c, a_1, a_2\}$ to a pair of pants decomposition P consisting of nonseparating circles on R in any way we want. Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c', a'_1, a'_2 be the representatives of $\lambda([a]), \lambda([b]), \lambda([c]), \lambda([a_1]), \lambda([a_2])$ in P' respectively. Since b is adjacent to a, c, a_1, a_2 w.r.t. P , b' is adjacent to a', c', a'_1, a'_2 w.r.t. P' . Then there are two pairs of pants Q_1, Q_2 having b' as boundary components and $Q_1 \cup Q_2$ has a', c', a'_1, a'_2 on its boundary. W.L.O.G. assume that Q_1 has a', b' on its boundary. Let x be the other boundary component of Q_1 . If $x = c'$ then we are done. Assume that $x \neq c'$. Then x is either a'_1 or a'_2 . Assume that $x = a'_1$. Since a_3 intersects b essentially and a_3 is disjoint from $a \cup a_1$, we can choose a representative a'_3 of $\lambda([a_3])$ such that there exists an essential arc w of a'_3 in Q_1 which starts and ends on b' and which does not intersect $a' \cup a'_1$. Now, we consider a_4 ; a_4 is disjoint from b and a_3 . Then there exists a representative a'_4 of $\lambda([a_4])$ such that a'_4 is disjoint from $b' \cup a'_3$. But then a'_4 could be isotoped so that it is disjoint from a' , since a' is a boundary component of a regular neighborhood of $b' \cup w$ in Q_1 . This is a contradiction since $i([a_4], [a]) \neq 0$ and so $i(\lambda([a_4]), \lambda([a])) \neq 0$. So, $x \neq a'_1$. Now, assume that $x = a'_2$. We consider the curves a_5, a_6 on N as shown in Figure 2 (iii). Since a_5 intersects c essentially and a_5 is disjoint from a_1, b , we can choose a representative a'_5 of $\lambda([a_5])$ such that there exists an essential arc w of a'_5 in Q_2 which starts and ends on c' and which does not intersect $a'_1 \cup b'$. The circle a_6 is disjoint from c and a_5 . Then there exists a representative a'_6 of $\lambda([a_6])$ such that a'_6 is disjoint from $c' \cup a'_5$. But then a'_6 could be isotoped so that it is disjoint from a'_1 , since a'_1 is a boundary component of a regular neighborhood of $c' \cup w$ in Q_2 . This is a contradiction since $i([a_6], [a_1]) \neq 0$ and so $i(\lambda([a_6]), \lambda([a_1])) \neq 0$. This completes the proof of the lemma. $\square$

Let α, β be two distinct vertices in $\mathcal{N}(R)$. We call (α, β) to be a *peripheral pair* in $\mathcal{N}(R)$ if they have disjoint representatives x, y respectively such that x, y and a boundary component of R bound a pair of pants in R .

Lemma 2.5 *Suppose $g \geq 2$ and $p \geq 1$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map and (α, β) be a peripheral pair in $\mathcal{N}(R)$. Then $(\lambda(\alpha), \lambda(\beta))$ is a peripheral pair in $\mathcal{N}(R)$.*

Proof. Let x, y be disjoint representatives of α, β respectively such that x, y and a boundary component of R bound a pair of pants in R . If $g = 2, p = 1$ then we complete x, y to a

pair of pants decomposition P consisting of nonseparating circles a, b on R . Then there are two distinct pair pants in P , one of them has a, b, x on its boundary and the other has a, b, y on its boundary. Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let x', y', a', b' be the representatives of $\lambda([x]), \lambda([y]), \lambda([a]), \lambda([b])$ in P' respectively. By the previous lemma, there exist two pairs of pants Q_1, Q_2 in P' such that Q_1 has a', b', x' on its boundary and Q_2 has a', b', y' on its boundary. Then it is clear that x', y' and the boundary component of R bound a pair of pants, which proves the lemma for this case. If $g \geq 3, p = 1$ then the proof is similar.

If $g = 2, p = 2$ then it is easy to see that we can complete x, y to a pair of pants decomposition P consisting of nonseparating circles a, b, z on R such that $([y], [z])$ is a peripheral pair, a, b, x bound a pair of pants in P and also a, b, z bound a pair of pants in P . Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let x', y', z', a', b' be the representatives of $\lambda([x]), \lambda([y]), \lambda([z]), \lambda([a]), \lambda([b])$ in P' respectively. By the previous lemma, there exist two pairs of pants Q_1, Q_2 in P' such that Q_1 has a', b', x' on its boundary and Q_2 has a', b', z' on its boundary. Then, since x is adjacent to y w.r.t. P , x' is adjacent to y' w.r.t. P' . So, there exists a pair of pants Q_3 in P' having x', y' on its boundary. Let w be the third boundary component of Q_3 . It is easy to see that $Q_1 \cup Q_2 \cup Q_3$ is a genus one surface with three boundary components, w, y', z' . If $w = z'$ then y' would be a separating curve which is a contradiction. If $w = y'$ then z' would be a separating curve which is a contradiction. Then it is clear that w has to be a boundary component of R , which proves the lemma for this case.

Assume that $g = 2, p \geq 3$. We choose distinct essential circles z, t, w as shown in Figure 3 (i). Then we complete x, y, z, t, w to a pair of pants decomposition P consisting of nonseparating circles in any way we like. Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let x', y', z', t', w' be the representatives of $\lambda([x]), \lambda([y]), \lambda([z]), \lambda([t]), \lambda([w])$ in P' respectively. Since z is a 4-curve in P , z' is a 4-curve in P' . Since x, z, w are the boundary components of a pair of pants in P , by the previous lemma, x', z', w' are the boundary components of a pair of pants, Q_1 , in P' . Similarly, since z, y, t are the boundary components of a pair of pants in P , by the previous lemma, z', y', t' are the boundary components of a pair of pants, Q_2 , in P' . Since x is adjacent to y in P , x' is adjacent to y' in P' . Then there exists a pair of pants Q_3 such that Q_3 has x', y' on its boundary. Let r be the boundary component of Q_3 different from x' and y' . Suppose that r is not a boundary component of R . Then r is an essential circle. Then $Q_1 \cup Q_2 \cup Q_3$ is a genus one subsurface with three boundary components r, w', t' which are nonseparating circles in R . Each two of x', w', t' can be connected by an arc in the complement of $Q_1 \cup Q_2 \cup Q_3$ in R . But this would be possible only if genus of R is at least 3. Since we assumed that $g = 2$, we get a contradiction. So, r has to be a boundary component of R , i.e. $(\lambda(\alpha), \lambda(\beta))$ is a peripheral pair in $\mathcal{N}(R)$.

Now, assume that $g = 3, p \geq 2$. We choose distinct essential circles $a_1, \dots, a_6$ as shown in Figure 3 (ii). Then we complete $x, y, a_1, \dots, a_6$ to a pair of pants decomposition P consisting of nonseparating circles in any way we like. Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let $x', y', a'_1, \dots, a'_6$ be the representatives of $\lambda([x]), \lambda([y]), \lambda([a_1]), \dots, \lambda([a_6])$ in P' respectively. Since a_1 is a 4-curve in P , a'_1 is a 4-curve in P' . Since x, a_1, a_3 are the boundary components of a pair of pants in P , by the previous lemma, x', a'_1, a'_3 are the boundary components of a pair of pants, Q_1 , in P' . Similarly, since y, a_1, a_2 are the boundary components of a pair of pants in P , by the previous lemma, y', a'_1, a'_2 are the boundary components of a pair of pants, Q_2 , in P' . By using similar arguments we can see that a'_4 is a 4-curve, a'_3, a'_4, a'_6 are the boundary components of a pair of pants, Q_3 , in P' and a'_2, a'_4, a'_5 are the boundary components of a pair of pants, Q_4 in




Figure 3: Pants decompositions

P' . Since x is adjacent to y in P , x' is adjacent to y' in P' . Then there exists a pair of pants Q_5 such that Q_5 has x', y' on its boundary. Let r be the boundary component of Q_5 different from x' and y' . Suppose that r is not a boundary component of R . Then r is an essential circle. Then $Q_1 \cup Q_2 \cup Q_3 \cup Q_4 \cup Q_5$ is a genus two subsurface with three boundary components r, a'_5, a'_6 which are nonseparating circles in R . Each two of r, a'_5, a'_6 can be connected by an arc in the complement of $Q_1 \cup Q_2 \cup Q_3 \cup Q_4 \cup Q_5$ in R . But this would be possible only if genus of R is at least 4. Since we assumed that $g = 3$, we get a contradiction. So, r has to be a boundary component of R , i.e. $(\lambda(\alpha), \lambda(\beta))$ is a peripheral pair in $\mathcal{N}(R)$. The proof of the case when $g \geq 4, p \geq 2$ is similar. $\square$

Lemma 2.6 *Suppose $g \geq 2$ and $p \geq 0$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Then λ preserves topological equivalence of ordered pairs of pants decompositions consisting of nonseparating circles on R , (i.e. for a given ordered pair of pants decomposition $P = (c_1, c_2, \dots, c_{3g-3+p})$ of R where $[c_i] \in \mathcal{N}(R)$, and a corresponding ordered pair of pants decomposition $P' = (c'_1, c'_2, \dots, c'_{3g-3+p})$ of R , where $[c'_i] = \lambda([c_i]) \forall i = 1, 2, \dots, 3g-3+p$, there exists a homeomorphism $H : R \rightarrow R$ such that $H(c_i) = c'_i \forall i = 1, 2, \dots, 3g-3+p$).*

Proof. Let $P = (c_1, c_2, \dots, c_{3g-3+p})$ be an ordered pair of pants decomposition consisting of nonseparating circles on R . Let $c'_i \in \lambda([c_i])$ such that the elements of $\{c'_1, c'_2, \dots, c'_{3g-3+p}\}$ are pairwise disjoint. Then $P' = (c'_1, c'_2, \dots, c'_{3g-3+p})$ is an ordered pair of pants decomposition of R . Let $(B_1, B_2, \dots, B_m)$ be an ordered set of all the pairs of pants in P . By Lemma 2.4 and Lemma 2.5, there is a corresponding, “image”, ordered collection of pairs of pants $(B'_1, B'_2, \dots, B'_m)$. A pair of pants having three essential curves on its boundary corresponds to a pair of pants having three essential curves on its boundary. A pair of pants having an inessential boundary component corresponds to a pair of pants having an inessential boundary component. Then by the classification of surfaces, there exists an orientation preserving homeomorphism $h_i : B_i \rightarrow B'_i$, for all $i = 1, \dots, m$. We can compose each h_i with an orientation preserving homeomorphism r_i which switches the boundary components, if necessary, to get $h'_i = r_i \circ h_i$ to agree with the correspondence given by λ on the boundary components, (i.e. for each essential boundary component a of B_i for $i = 1, \dots, m$, $\lambda([q(a)]) = [q'(h'_i(a))]$ where $q : R_P \rightarrow R$ and $q' : R_{P'} \rightarrow R$ are the natural quotient maps). Then for two pairs of pants with a common boundary component, we can glue the homeomorphisms by isotoping the homeomorphism of the one pair of pants so that it agrees with the homeomorphism of the other pair of pants on the common boundary component. By adjusting these homeomorphisms on the boundary components and gluing them we get a homeomorphism $h : R \rightarrow R$ such that $h(c_i) = c'_i$ for all $i = 1, 2, \dots, 3g-3+p$. $\square$

Lemma 2.7 *Suppose $g \geq 2$ and $p \geq 0$. Let α_1 and α_2 be two vertices in $\mathcal{N}(R)$. Then $i(\alpha_1, \alpha_2) = 1$ if and only if there exist isotopy classes $\alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7$ in $\mathcal{N}(R)$ such that*




Figure 4: Circles intersecting once

(i) $i(\alpha_i, \alpha_j) = 0$ if and only if the i^{th} and j^{th} circles on Figure 4 are disjoint.

(ii) $\alpha_1, \alpha_3, \alpha_5, \alpha_6$ have pairwise disjoint representatives a_1, a_3, a_5, a_6 respectively such that $a_5 \cup a_6$ divides R into two pieces, one of these is a torus with two holes, T , containing some representatives of the isotopy classes α_1, α_2 and a_1, a_3, a_5 bound a pair of pants in T and a_1, a_3, a_6 bound a pair of pants in T .

Proof. Let $i(\alpha_1, \alpha_2) = 1$. Let a_1, a_2 be representatives of α_1, α_2 respectively such that a_1 intersects a_2 transversely once. Let N be a regular neighborhood of $a_1 \cup a_2$. Then it is easy to see that N is a genus one surface with one boundary component and since R is a surface of genus at least 2, there are simple closed curves as shown in Figure 4 such that their isotopy classes $\alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7$ satisfy the properties (i) and (ii). Suppose that there exist isotopy classes $\alpha_1, \alpha_2, \dots, \alpha_7$ in $\mathcal{N}(R)$ satisfying (i) and (ii). Then we have a_1, a_3, a_5, a_6 as pairwise disjoint representatives of $\alpha_1, \alpha_3, \alpha_5, \alpha_6$ respectively such that $a_5 \cup a_6$ divides R into two pieces, one of these is a torus with two holes, T , containing some representatives of the isotopy classes α_1, α_2 and a_1, a_3, a_5 bound a pair of pants P in T , and a_1, a_3, a_6 bound a pair of pants Q in T . Let a_2, a_4, a_7 be representatives of $\alpha_2, \alpha_4, \alpha_7$ such that all the curves a_i , $i = 1, \dots, 7$ have minimal intersection with each other. Then we have $a_4 \cap a_1 = \emptyset, a_2 \cap a_6 = \emptyset, a_7 \cap a_3 = \emptyset$. Since $i(\alpha_4, \alpha_1) = 0$, $i(\alpha_4, \alpha_3) \neq 0$ and $i(\alpha_4, \alpha_6) \neq 0$, $a_4 \cap Q$ has an arc which connects a_3 to a_6 . Since $i(\alpha_7, \alpha_3) = 0$, $i(\alpha_7, \alpha_1) \neq 0$ and $i(\alpha_7, \alpha_6) \neq 0$, $a_7 \cap Q$ has an arc which connects a_1 to a_6 . Similarly, since $i(\alpha_2, \alpha_6) = 0$, $i(\alpha_2, \alpha_1) \neq 0$ and $i(\alpha_2, \alpha_3) \neq 0$, $a_2 \cap Q$ has an arc which connects a_1 to a_3 . Then, since a_2, a_4, a_7 are pairwise disjoint, we can see that all the arcs of $a_4 \cap Q$ connect a_3 to a_6 , all the arcs of $a_7 \cap Q$ connect a_1 to a_6 and all the arcs of $a_2 \cap Q$ connect a_1 to a_3 . By using similar arguments, we can see that all the arcs of $a_4 \cap P$ connect a_3 to a_5 , all the arcs of $a_7 \cap P$ connect a_1 to a_5 , and all the arcs of $a_2 \cap P$ connect a_1 to a_3 . Then by looking at the gluing between the arcs in $a_2 \cap Q$ and the arcs in $a_2 \cap P$ to form a_2 , we see that $a_2 \cap Q$ has one arc and $a_2 \cap P$ has one arc. Hence, $i(\alpha_1, \alpha_2) = 1$. $\square$

A characterization of geometric intersection one property in $\mathcal{C}(R)$ was given by Ivanov, in Lemma 1 in [5].

Lemma 2.8 *Suppose $g \geq 2$ and $p \geq 0$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Let α, β be two vertices of $\mathcal{N}(R)$. If $i(\alpha, \beta) = 1$, then $i(\lambda(\alpha), \lambda(\beta)) = 1$.*

Proof. Let α, β be two vertices of $\mathcal{N}(R)$ such that $i(\alpha, \beta) = 1$. Then by Lemma 2.7, there exist isotopy classes $\alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7$ in $\mathcal{N}(R)$ such that $i(\alpha_i, \alpha_j) = 0$ if and only if $i^{\text{th}}, j^{\text{th}}$ circles on Figure 4 are disjoint and $\alpha_1, \alpha_3, \alpha_5, \alpha_6$ have pairwise disjoint representatives

a_1, a_3, a_5, a_6 respectively such that $a_5 \cup a_6$ divides R into two pieces, one of these is a torus with two holes, T , containing some representatives of the isotopy classes α_1, α_2 and a_1, a_3, a_5 bound a pair of pants in T and a_1, a_3, a_6 bound a pair of pants in T . Then, since λ is superinjective, $i(\lambda(\alpha_i), \lambda(\alpha_j)) = 0$ if and only if i^{th}, j^{th} circles on Figure 4 are disjoint, and by using Lemma 2.6 and the properties that λ preserves disjointness and nondisjointness, we can see that there are pairwise disjoint representatives a'_1, a'_3, a'_5, a'_6 of $\lambda(\alpha_1), \lambda(\alpha_3), \lambda(\alpha_5), \lambda(\alpha_6)$ respectively, such that $a'_5 \cup a'_6$ divides R into two pieces, one of these is a torus with two holes, T , containing some representatives of the isotopy classes $\lambda(\alpha_1), \lambda(\alpha_2)$ and a'_1, a'_3, a'_5 bound a pair of pants in T and a'_1, a'_3, a'_6 bound a pair of pants in T . Then by Lemma 2.7, $i(\lambda(\alpha), \lambda(\beta)) = 1$. $\square$

3 Superinjective Simplicial Maps of $\mathcal{C}(R)$ and Injective Homomorphisms of Finite Index Subgroups of Mod_R^*

If $g = 2, p \geq 2$ or $g \geq 3, p \geq 0$, by the results given in [3] and [4], we have the following: A simplicial map $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is superinjective if and only if λ is induced by a homeomorphism of R . If K is a finite index subgroup of Mod_R^* and $f : K \rightarrow Mod_R^*$ is an injective homomorphism, then f is induced by a homeomorphism of R . In this section we prove similar results, Theorem 1.3 and Theorem 1.4, when $g = 2, p \leq 1$.

If $g = 2$ and $p \leq 1$, and $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is a superinjective simplicial map, then λ is an injective simplicial map which maps pair of pants decompositions of R to pair of pants decompositions of R and it preserves adjacency relation. The proofs of these are similar to the proofs given in Lemma 1.1-1.3. Now, we prove the following lemma:

Lemma 3.1 *Suppose $g = 2$ and $p \leq 1$. Let $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ be a superinjective simplicial map and α, β, γ be distinct vertices in $\mathcal{C}(R)$ having pairwise disjoint representatives which bound a pair of pants in R . Then $\lambda(\alpha), \lambda(\beta), \lambda(\gamma)$ are distinct vertices in $\mathcal{C}(R)$ having pairwise disjoint representatives which bound a pair of pants in R .*

Proof. Let a, b, c be pairwise disjoint representatives of α, β, γ respectively. If R is a closed surface of genus two then a, b, c is a pair of pants decomposition on R and since a, b, c bound a pair of pants, they have to be all nonseparating circles in this case. Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c' be the representatives of $\lambda([a]), \lambda([b]), \lambda([c])$ in P' respectively. Since a is adjacent to b and c w.r.t. P , a' is adjacent to b' and c' w.r.t. P' . If one of a', b' or c' was a separating curve on R then the other two wouldn't be adjacent to each other w.r.t. P' , which would give a contradiction. So, each of a', b', c' is a nonseparating curve. Then clearly they bound a pair of pants on R . Now assume that $g = 2, p = 1$. There are two cases: either each of a, b, c is a nonseparating circle or exactly one of a, b, c is a separating circle.

Case i: Assume that each of a, b, c is a nonseparating circle and complete $\{a, b, c\}$ to a pair of pants decomposition $P = \{a, b, c, a_1\}$ consisting of nonseparating circles as shown in Figure 5 (i). Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c', a'_1 be the representatives of $\lambda([a]), \lambda([b]), \lambda([c]), \lambda([a_1])$ in P' respectively. Notice that any two curves in P are adjacent w.r.t. P . Since adjacency is preserved, any two curves in P' must be adjacent w.r.t. P' . If one of a', b', c', a'_1 was a separating curve then there would be two circles in P' which are not adjacent. So, we conclude that all of a', b', c', a'_1 are nonseparating. Then we consider the curve configuration as shown in Figure 5 (i) and the proof of the lemma in this case follows as in the proof of Lemma 2.4 for $g = 2, p = 1$ case.




Figure 5: Curve configurations

Case ii: Assume that exactly one of a, b, c is a separating circle. W.L.O.G. assume that c is a separating circle. Then c separates R into two subsurfaces R_1, R_2 as shown in Figure 5 (ii). Let a_1, a_2, a_3 be as shown in the figure. Then a, b, c, a_1 is a pants decomposition P on R . Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c', a'_1 be the representatives of $\lambda([a]), \lambda([b]), \lambda([c]), \lambda([a_1])$ in P' respectively. Every nonseparating circle x on R could be put inside of a pair of pants decomposition consisting of nonseparating circles, and using the method given in case (i), we could see that $\lambda([x])$ has a nonseparating representative. So, since a, b, a_1 are nonseparating, a', b', a'_1 are nonseparating. Since a is adjacent to c w.r.t. P , a' is adjacent to c' w.r.t. P' . Then there is a pair of pants Q_1 having a' and c' on its boundary. Let x be the other boundary component of Q_1 . Since a' is not a separating circle, x can't be c' . If x is the boundary component of R , then since c' is adjacent to a'_1 and b' , there exists a pair of pants Q_2 in P' having c', b' and a'_1 on its boundary. Then, since a_2 intersects a_1 essentially and a_2 is disjoint from b and c , we can choose a representative a'_2 of $\lambda([a_2])$ such that there exists an essential arc w of a'_2 in Q_2 which starts and ends on a'_1 and which does not intersect $b' \cup c'$. Now, we consider a_3 which is disjoint from a_1 and a_2 . Then there exists a representative a'_3 of $\lambda([a_3])$ such that a'_3 is disjoint from $a'_1 \cup w$. But then a'_3 could be isotoped so that it is disjoint from b' , since b' is a boundary component of a regular neighborhood of $a'_1 \cup w$ in Q_2 . This is a contradiction since $i([b], [a_3]) \neq 0$ and so $i(\lambda([b]), \lambda([a_3])) \neq 0$. So, x can not be the boundary component of R . Then there are three possibilities: x is one of a', b', a'_1 . If $x = a'$ then a' wouldn't be adjacent to b' w.r.t. P' which is a contradiction. Assume $x = a'_1$. Then, since a_2 intersects a_1 essentially and a_2 is disjoint from a and c , we can choose a representative a'_2 of $\lambda([a_2])$ such that there exists an essential arc w of a'_2 in Q_1 which starts and ends on a'_1 and which does not intersect $a' \cup c'$. Now, we consider a_3 which is disjoint from a_1 and a_2 . Then there exists a representative a'_3 of $\lambda([a_3])$ such that a'_3 is disjoint from $a'_1 \cup w$. But then a'_3 could be isotoped so that it is disjoint from a' , since a' is a boundary component of a regular neighborhood of $a'_1 \cup w$ in Q_1 . This is a contradiction since $i([a], [a_3]) \neq 0$ and so $i(\lambda([a]), \lambda([a_3])) \neq 0$. So, $x \neq a'_1$. Then $x = b'$, and a', b', c' bound a pair of pants in P' . $\square$

Let α, β be two distinct vertices in $\mathcal{C}(R)$. We call (α, β) to be a *peripheral pair* in $\mathcal{C}(R)$ if they have disjoint representatives x, y respectively such that x, y and a boundary component of R bound a pair of pants in R .

Lemma 3.2 Suppose $g = 2$ and $p = 1$. Let $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ be a superinjective simplicial map and (α, β) be a peripheral pair in $\mathcal{C}(R)$. Then $(\lambda(\alpha), \lambda(\beta))$ is a peripheral pair in $\mathcal{C}(R)$.

Proof. Let x, y be disjoint representatives of α, β respectively such that x, y and a boundary component of R bound a pair of pants in R . There are two cases to consider: Case i: x and y are both nonseparating. Case ii: x and y are both separating. The proof in the first

case is similar to the proof given in Lemma 2.5. For the second case, we complete x, y to a pair of pants decomposition Q consisting of nonseparating circles a, b on R such that a is in the torus with one hole which comes with the separation by x . Then we will replace y with a nonseparating curve w such that a, x, b, w is a pants decomposition P on R , x, w, b bound a pair of pants and w, b and the boundary component of R bound a pair of pants on R . Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let x', w', a', b' be the representatives of $\lambda([x]), \lambda([w]), \lambda([a]), \lambda([b])$ in P' respectively. Then by using case i and Lemma 3.1, we see that x' is a separating circle of genus one. Similarly, y' is a separating circle of genus one on R . Then it is easy to see that x', y' and the boundary component of R bound a pair of pants. $\square$

Lemma 3.3 *Suppose $g = 2$ and $p \leq 1$. Let $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ be a superinjective simplicial map. Then λ preserves topological equivalence of ordered pairs of pants decompositions on R , (i.e. for a given ordered pair of pants decomposition $P = (c_1, c_2, \dots, c_{3+p})$ of R where $[c_i] \in \mathcal{C}(R)$, and a corresponding ordered pair of pants decomposition $P' = (c'_1, c'_2, \dots, c'_{3+p})$ of R , where $[c'_i] = \lambda([c_i]) \forall i = 1, 2, \dots, 3+p$, there exists a homeomorphism $H : R \rightarrow R$ such that $H(c_i) = c'_i \forall i = 1, 2, \dots, 3+p$).*

Proof. First we will consider the case when $g = 2$ and $p = 0$. Let P be a pair of pants decomposition of R and A be a nonembedded pair of pants in P . The boundary of A consists of the circles a, c where c is a 1-separating circle on R and a is a nonseparating circle on R . Let b, a_1, a_2 be as shown in Figure 5 (iii). Then $P = \{a, b, c\}$ is a pants decomposition on R . Let P' be a pair of pants decomposition of R such that $\lambda([P]) = [P']$. Let a', b', c' be the representatives of $\lambda([a]), \lambda([b]), \lambda([c])$ in P' respectively. Every nonseparating circle x on R could be put inside of a pair of pants decomposition consisting of nonseparating circles, and using the method given in case (i) in Lemma 3.1, we could see that $\lambda([x])$ has a nonseparating representative. So, since a, b are nonseparating, a', b' are nonseparating. Assume that c' is also nonseparating. Then there exist two pairs of pants Q_1, Q_2 in P' such that both of them have a', b', c' on their boundary. Then, since a_1 intersects a essentially and a_1 is disjoint from b and c , we can choose a representative a'_1 of $\lambda([a_1])$ such that there exists an essential arc w of a'_1 in Q_1 which starts and ends on a' and which does not intersect $b' \cup c'$. Now, we consider a_2 which is disjoint from a and a_1 . There exists a representative a'_2 of $\lambda([a_2])$ such that a'_2 is disjoint from $a'_1 \cup a'$. But then a'_2 could be isotoped so that it is disjoint from b' , since b' is a boundary component of a regular neighborhood of $a' \cup w$ in Q_1 . This is a contradiction since $i([b], [a_2]) \neq 0$ and so $i(\lambda([b]), \lambda([a_2])) \neq 0$. So, c' has to be separating. Then clearly a' and c' are the boundary components of a nonembedded pair of pants. Hence, nonembedded pair of pants in P corresponds to a nonembedded pair of pants in P' . When $g = 2$ and $p = 1$, this result follows from Lemma 3.1 and Lemma 3.2 considering the circles in Figure 5 (ii).

Suppose $g = 2$ and $p \leq 1$. Let $P = (c_1, c_2, \dots, c_{3+p})$ be an ordered pair of pants decomposition on R . Let $c'_i \in \lambda([c_i])$ such that the elements of $\{c'_1, c'_2, \dots, c'_{3+p}\}$ are pairwise disjoint. Then $P' = (c'_1, c'_2, \dots, c'_{3+p})$ is an ordered pair of pants decomposition of R . Let $(B_1, B_2, \dots, B_m)$ be an ordered set of all the pairs of pants in P . By Lemma 3.1, Lemma 3.2 and the arguments given above, there is a corresponding, “image”, ordered collection of pairs of pants $(B'_1, B'_2, \dots, B'_m)$. Nonembedded pairs of pants correspond to nonembedded pairs of pants, embedded pairs of pants correspond to embedded pairs of pants, and a pair of pants having an inessential boundary component corresponds to a pair of pants having an inessential boundary component. Then the proof follows as in Lemma 2.6. $\square$

Lemma 3.4 *Suppose $g = 2$ and $p \leq 1$. Let $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ be a superinjective simplicial map. Let α, β be two vertices of $\mathcal{C}(R)$. If $i(\alpha, \beta) = 1$, then $i(\lambda(\alpha), \lambda(\beta)) = 1$.*




Figure 6: Curves intersecting twice

Proof. The proof follows as in the proof of Lemma 2.8, using Lemma 3.3. $\square$

Lemma 3.5 *Suppose $g = 2$ and $p \leq 1$. Let $\lambda : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ be a superinjective simplicial map. Let α and β be two vertices in $\mathcal{C}(R)$ which have representatives with geometric intersection 2 and algebraic intersection 0 on R . Then $\lambda(\alpha)$ and $\lambda(\beta)$ have representatives with geometric intersection 2 and algebraic intersection 0 on R .*

Proof. Assume that $g = 2$ and $p = 0$. Let h, v be representatives of α, β with geometric intersection 2 and algebraic intersection 0 on R . Let N be a regular neighborhood of $h \cup v$ in R . Then N is a sphere with four boundary components. Let c, x, y, z be boundary components of N such that there exists a homeomorphism $\varphi : (N, c, x, y, z, h, v) \rightarrow (N_o, c_o, x_o, y_o, z_o, h_o, v_o)$ where N_o is a standard sphere with four holes having c_o, x_o, y_o, z_o on its boundary and h_o, v_o (horizontal, vertical) are two circles as indicated in Figure 6 (i). Since h and v have geometric intersection 2 and algebraic intersection 0 on R , none of c, x, y, z bound a disk on R . There are two cases to consider: either exactly one of h or v is separating or both h and v are nonseparating. Case i: W.L.O.G. Assume that v is separating and h is nonseparating. Then x is isotopic to z and y is isotopic to c and c, x, h, v are as shown in Figure 6 (ii). Let a_1, a_2 be as shown in the figure. Let c', h', x' be pairwise disjoint representatives of $\lambda([c]), \lambda([h]), \lambda([x])$ respectively. Then $\{c', h', x'\}$ is a pair of pants decomposition on R . Since nondisjointness and intersection one is preserved by λ , there are disjoint representatives a'_1, a'_2 of $\lambda([a_1]), \lambda([a_2])$ respectively such that a'_1 intersects each of c' and h' exactly once and a'_1 is disjoint from x' , and similarly a'_2 intersects each of x' and h' exactly once and a'_2 is disjoint from c' . Then, since v is disjoint from $c \cup a_1 \cup a_2 \cup x$, we can choose a representative v' of $\lambda([v])$ such that it lives inside of the cylinder that we get after cutting R along $c' \cup a'_1 \cup a'_2 \cup x'$. Then it is easy to see that h' and v' intersects geometrically twice and algebraically zero times.

Case ii: Assume that both of h, v are nonseparating. Then c and z are the boundary components of an annulus, A_1 , in $R \setminus N$ and x and y are the boundary components of an annulus, A_2 , in $R \setminus N$. Let i and j be as shown in Figure 7 (i). We connect the ends points of i on N with an arc in A_1 to get a circle w such that w intersects each of c and z at only one point. Similarly, we connect the ends points of j on N with an arc in A_2 to get a circle k such that k intersects each of x and y at only one point. Notice that w and k intersect at only one point. Let c', x', h', y', z' be pairwise disjoint representatives of the image curves. Let N' be sphere with four holes bounded by c', x', y', z' . By Lemma 3.3, h' gives a pants decompositions on N' . Now we choose minimally intersecting representatives w', k' of $\lambda([w]), \lambda([k])$ respectively such that each of w' and k' intersects c', x', y', z' minimally. Then, since the intersection one is preserved, there are arcs i', j' of w', k' in N'




Figure 7: Curves intersecting twice

respectively such that i' intersects j' once in N' , i' connects c' to z' and j' connects x' to y' . Since v does not intersect any of x, c, y, z , and v intersects w and k exactly once, there is a representative v' of $\lambda([v])$ in N' such that v' intersects each of i' and j' exactly once. Notice that h' also intersects each of i' and j' exactly once. When we cut N' along the arcs of i' and j' we get a disk. The boundary of the disk either is as shown in Figure 7 (ii) or it has the similar form where only x' and y' are switched. We will consider the first case (the arguments follow in the second case similarly). If v' makes its intersection with i' and j' at the intersection point of i' and j' with each other, then v' has to be one of the arcs shown in Figure 7 (ii), and by looking at the intersection of v' with the other curves, it is easy to see that v' intersects h' geometrically twice and algebraically 0 times on R . Suppose that v' intersects i' and j' at different points. Then there are two arcs of v' connecting i' to j' in the disk. Then, since v' does not intersect any of x', c', y', z' we see that v' has to be one of the curves as shown in Figure 7 (iii), (iv). Since v' is not isotopic to h' , v' has to be the curve shown in Figure 7 (iii) and hence v' intersects h' geometrically twice and algebraically 0 times on R . If $g = 2$ and $p = 1$ the proof is similar (for Case (i) we use Figure 6 (iii)). $\square$

The proof of Theorem 1.3 follows from Lemma 3.3, Lemma 3.4, Lemma 3.5, and the techniques given for the proof of Theorem 1.1 in [3], [4].

A mapping class $g \in \text{Mod}_R^*$ is called *pseudo-Anosov* if $\mathcal{A}$ is nonempty and if $g^n(\alpha) \neq \alpha$, for all α in $\mathcal{A}$ and any $n \neq 0$. g is called *reducible* if there is a nonempty subset $\mathcal{B} \subseteq \mathcal{A}$ such that a set of disjoint representatives can be chosen for $\mathcal{B}$ and $g(\mathcal{B}) = \mathcal{B}$. In this case, $\mathcal{B}$ is called a *reduction system* for g . Each element of $\mathcal{B}$ is called a *reduction class* for g . A reduction class, α , for g , is called an *essential reduction class* for g , if for each $\beta \in \mathcal{A}$ such that $i(\alpha, \beta) \neq 0$ and for each integer $m \neq 0$, $g^m(\beta) \neq \beta$. The set, $\mathcal{B}_g$, of all essential reduction classes for g is called the *canonical reduction system* for g .

Let $\Gamma' = \ker(\varphi)$ where $\varphi : \text{Mod}_R^* \rightarrow \text{Aut}(H_1(R, \mathbb{Z}_3))$ is the homomorphism defined by the action of homeomorphisms on the homology. The proofs of the Lemma 3.6 and Lemma 3.7 follow by the techniques given in [3]. Note that we need to use that the maximal rank of an abelian subgroup of Mod_R^* is $3g - 3 + p$, [1].

Lemma 3.6 *Suppose $g = 2$ and $p \leq 1$. Let K be a finite index subgroup of Mod_R^* and $f : K \rightarrow \text{Mod}_R^*$ be an injective homomorphism. Let $\alpha \in \mathcal{A}$. Then there exists $N \in \mathbb{Z}^*$ such that $\text{rank } C(C_{\Gamma'}(f(t_\alpha^N))) = 1$.*

Lemma 3.7 Suppose $g = 2$ and $p \leq 1$. Let K be a finite index subgroup of Mod_R^* . Let $f : K \rightarrow \text{Mod}_R^*$ be an injective homomorphism. Then there exists $N \in \mathbb{Z}^*$ such that $f(t_\alpha^N)$ is a reducible element of infinite order for all $\alpha \in \mathcal{A}$.

In the proof of Lemma 3.7, we use that the centralizer of a p-Anosov element in the extended mapping class group is a virtually infinite cyclic group, [10].

Lemma 3.8 Suppose $g = 2$ and $p \leq 1$. Let K be a finite index subgroup of Mod_R^* and $f : K \rightarrow \text{Mod}_R^*$ be an injective homomorphism. Then $\forall \alpha \in \mathcal{A}$, $f(t_\alpha^N) = t_{\beta(\alpha)}^M$ for some $M, N \in \mathbb{Z}^*$, $\beta(\alpha) \in \mathcal{A}$.

Proof. Let $\Gamma = f^{-1}(\Gamma') \cap \Gamma'$. Since Γ is a finite index subgroup we can choose $N \in \mathbb{Z}^*$ such that $t_\alpha^N \in \Gamma$ for all α in $\mathcal{A}$. By Lemma 3.7 $f(t_\alpha^N)$ is a reducible element of infinite order in Mod_R^* . Let C be a realization of the canonical reduction system of $f(t_\alpha^N)$. Let c be the number of components of C and r be the number of p-Anosov components of $f(t_\alpha^N)$. Since $t_\alpha^N \in \Gamma$, $f(t_\alpha^N) \in \Gamma'$. By Theorem 5.9 [7], $C(C_{\Gamma'}(f(t_\alpha^N)))$ is a free abelian group of rank $c + r$. By Lemma 3.6 $c + r = 1$. Then either $c = 1$, $r = 0$ or $c = 0$, $r = 1$. Since there is at least one curve in the canonical reduction system we have $c = 1$, $r = 0$. Hence, since $f(t_\alpha^N) \in \Gamma'$, $f(t_\alpha^N) = t_{\beta(\alpha)}^M$ for some $M \in \mathbb{Z}^*$, $\beta(\alpha) \in \mathcal{A}$, [1], [7]. $\square$

Remark: Suppose that $f(t_\alpha^M) = t_\beta^P$ for some $\beta \in \mathcal{A}$ and $M, P \in \mathbb{Z}^*$ and $f(t_\alpha^N) = t_\gamma^Q$ for some $\gamma \in \mathcal{A}$ and $N, Q \in \mathbb{Z}^*$. Since $f(t_\alpha^{M \cdot N}) = f(t_\alpha^{N \cdot M})$, $t_\beta^P = t_\gamma^Q$, $P, Q, M, N \in \mathbb{Z}^*$. Then $\beta = \gamma$. Therefore, by Lemma 3.8, f gives a correspondence between isotopy classes of circles and f induces a map, $f_* : \mathcal{A} \rightarrow \mathcal{A}$, where $f_*(\alpha) = \beta(\alpha)$.

In the following lemma we use a well known fact that $ft_\alpha f^{-1} = t_{f(\alpha)}^{\epsilon(f)}$ for all α in $\mathcal{A}$, $f \in \text{Mod}_R^*$, where $\epsilon(f) = 1$ if f has an orientation preserving representative and $\epsilon(f) = -1$ if f has an orientation reversing representative.

Lemma 3.9 Let K be a finite index subgroup of Mod_R^* . Let $f : K \rightarrow \text{Mod}_R^*$ be an injective homomorphism. Assume that there exists $N \in \mathbb{Z}^*$ such that $\forall \alpha \in \mathcal{A}$, $\exists Q \in \mathbb{Z}^*$ such that $f(t_\alpha^N) = t_\alpha^Q$. If $g = 2$ and $p = 1$ then f is the identity on K . If $g = 2$ and $p = 0$ then $f(k) = ki^{m(k)}$ where i is the hyperelliptic involution on R and $m(k) \in \{0, 1\}$.

Proof. We use Ivanov's trick to see that $f(kt_\alpha^N k^{-1}) = f(t_{k(\alpha)}^{\epsilon(k) \cdot N}) = t_{k(\alpha)}^{Q \cdot \epsilon(k)}$ and $f(kt_\alpha^N k^{-1}) = f(k)f(t_\alpha^N)f(k)^{-1} = f(k)t_\alpha^Q f(k)^{-1} = t_{f(k)(\alpha)}^{\epsilon(f(k)) \cdot Q} \forall \alpha \in \mathcal{A}, \forall k \in K$. Then we have $t_{k(\alpha)}^{Q \cdot \epsilon(k)} = t_{f(k)(\alpha)}^{\epsilon(f(k)) \cdot Q} \forall \alpha \in \mathcal{A}, \forall k \in K$. Hence, $k(\alpha) = f(k)(\alpha) \forall \alpha \in \mathcal{A}, \forall k \in K$. Then $k^{-1}f(k)(\alpha) = \alpha \forall \alpha \in \mathcal{A}, \forall k \in K$. Therefore, $k^{-1}f(k)$ commutes with $t_\alpha \forall \alpha \in \mathcal{A}, \forall k \in K$. Then, since Mod_R is generated by Dehn twists when $g = 2, p \leq 1$, $k^{-1}f(k) \in C(\text{Mod}_R) \forall k \in K$. If $g = 2$ and $p = 1$, then $C(\text{Mod}_R)$ is trivial by 5.3 in [7]. So, $k = f(k) \forall k \in K$. Hence, $f = \text{id}_K$. If $g = 2$ and $p = 0$, then $C(\text{Mod}_R) = \{\text{id}_R, i\} = \mathbb{Z}_2$, where i is the hyperelliptic involution on R . Then for each $k \in K$ either $k^{-1}f(k) = \text{id}_R$ or $k^{-1}f(k) = i$. So, $f(k) = ki^{m(k)}$ where $m(k) \in \{0, 1\}$. $\square$

Corollary 3.10 Suppose $g = 2$ and $p = 1$. Let $h : \text{Mod}_R^* \rightarrow \text{Mod}_R^*$ be an isomorphism and $f : \text{Mod}_R^* \rightarrow \text{Mod}_R^*$ be an injective homomorphism. Assume that there exists $N \in \mathbb{Z}^*$ such that $\forall \alpha \in \mathcal{A}$, $\exists Q \in \mathbb{Z}^*$ such that $h(t_\alpha^N) = f(t_\alpha^Q)$. Then $h = f$.

Proof. Apply Lemma 3.9 to $h^{-1}f$ with $K = \text{Mod}_R^*$. Since for all α in $\mathcal{A}$, $h^{-1}f(t_\alpha^N) = t_\alpha^Q$, we have $h^{-1}f = \text{id}_K$. Hence, $h = f$. $\square$

By the remark after Lemma 3.8, we have that $f : K \rightarrow \text{Mod}_R^*$ induces a map $f_* : \mathcal{A} \rightarrow \mathcal{A}$, where K is a finite index subgroup of Mod_R^* . In the following lemma we prove that f_* is a superinjective simplicial map on $\mathcal{C}(R)$.

Lemma 3.11 *Suppose $g = 2$ and $p \leq 1$. Let $f : K \rightarrow \text{Mod}_R^*$ be an injection. Let $\alpha, \beta \in \mathcal{A}$. Then $i(\alpha, \beta) = 0 \Leftrightarrow i(f_*(\alpha), f_*(\beta)) = 0$.*

Proof. There exists $N \in \mathbb{Z}^*$ such that $t_\alpha^N \in K$ and $t_\beta^N \in K$. Then we have the following: $i(\alpha, \beta) = 0 \Leftrightarrow t_\alpha^N t_\beta^N = t_\beta^N t_\alpha^N \Leftrightarrow f(t_\alpha^N) f(t_\beta^N) = f(t_\beta^N) f(t_\alpha^N)$ (since f is injective on K) $\Leftrightarrow t_{f_*(\alpha)}^P t_{f_*(\beta)}^Q = t_{f_*(\beta)}^Q t_{f_*(\alpha)}^P$ where $P = M(\alpha, N), Q = M(\beta, N) \in \mathbb{Z}^* \Leftrightarrow i(f_*(\alpha), f_*(\beta)) = 0$. $\square$

Now, we prove the second main theorem of the section.

Theorem 3.12 *Let K be a finite index subgroup of Mod_R^* and f be an injective homomorphism $f : K \rightarrow \text{Mod}_R^*$. If $g = 2$ and $p = 1$ then f has the form $k \rightarrow hkh^{-1}$ for some $h \in \text{Mod}_R^*$ and f has a unique extension to an automorphism of Mod_R^* . If R is a closed surface of genus 2, then f has the form $k \rightarrow hkh^{-1}i^{m(k)}$ for some $h \in \text{Mod}_R^*$ where m is a homomorphism $K \rightarrow \mathbb{Z}_2$ and i is the hyperelliptic involution on R .*

Proof. If $g = 2$ and $p \leq 1$, by Lemma 3.11 f_* is a superinjective simplicial map on $\mathcal{C}(R)$. Then by Theorem 1.3 f_* is induced by a homeomorphism $h : R \rightarrow R$, i.e. $f_*(\alpha) = h_\#(\alpha)$ for all α in $\mathcal{A}$, where $h_\# = [h]$. Let $\chi^{h_\#} : \text{Mod}_R^* \rightarrow \text{Mod}_R^*$ be the isomorphism defined by the rule $\chi^{h_\#}(k) = h_\# k h_\#^{-1}$ for all k in Mod_R^* . Then for all α in $\mathcal{A}$, we have the following:

$$\chi^{h_\#^{-1}} \circ f(t_\alpha^N) = \chi^{h_\#^{-1}}(t_{f_*(\alpha)}^M) = \chi^{h_\#^{-1}}(t_{h_\#(\alpha)}^M) = h_\#^{-1} t_{h_\#(\alpha)}^M h_\# = t_{h_\#^{-1}(h_\#(\alpha))}^{M \cdot \epsilon(h_\#^{-1})} = t_\alpha^{M \cdot \epsilon(h_\#^{-1})}.$$

If $g = 2$ and $p = 1$, then since $\chi^{h_\#^{-1}} \circ f$ is injective, $\chi^{h_\#^{-1}} \circ f = \text{id}_K$ by Lemma 3.9. So, $\chi_\#^h|_K = f$. Hence, f is the restriction of an isomorphism which is conjugation by $h_\#$, (i.e. f is induced by h). Suppose that there exists an automorphism $\tau : \text{Mod}_R^* \rightarrow \text{Mod}_R^*$ such that $\tau|_K = f$. Let $N \in \mathbb{Z}^*$ such that $t_\alpha^N \in K$ for all α in $\mathcal{A}$. Since $\chi_\#^h|_K = f = \tau|_K$ and $t_\alpha^N \in K$, $\tau(t_\alpha^N) = \chi_\#^h(t_\alpha^N)$ for all α in $\mathcal{A}$. Then by Corollary 3.10, $\tau = \chi^{h_\#}$. Hence, the extension of f is unique.

If $g = 2$ and $p = 0$, then since $\chi^{h_\#^{-1}} \circ f$ is injective, $\chi^{h_\#^{-1}} \circ f(k) = ki^{m(k)}$ where $m(k) \in \{0, 1\}$ by Lemma 3.9. So, f has the form $k \rightarrow h_\# k h_\#^{-1} i^{m(k)}$. Since $\chi^{h_\#^{-1}} \circ f(k_1 k_2) = k_1 k_2 i^{m(k_1 k_2)}$ and $\chi^{h_\#^{-1}} \circ f(k_1) \chi^{h_\#^{-1}} \circ f(k_2) = k_1 i^{m(k_1)} k_2 i^{m(k_2)} = k_1 k_2 i^{m(k_1)} i^{m(k_2)}$, we have that $m(k_1 k_2) = m(k_1) + m(k_2)$ for all $k_1, k_2 \in K$. So, $m : K \rightarrow \mathbb{Z}_2$ is a homomorphism. $\square$

Remark: Note that $k \rightarrow hkh^{-1}i^{m(k)}$ defines a homomorphism from $K \rightarrow \text{Mod}_R^*$ for every $h \in \text{Mod}_R^*$ and for every homomorphism $m : K \rightarrow \mathbb{Z}_2$. It is easy to see that $k \rightarrow hkh^{-1}i^{m(k)}$ is injective if and only if either $i \notin K$ or $i \in \text{Ker}(m)$. Inner automorphisms of K act on the set of injective homomorphisms from $K \rightarrow \text{Mod}_R^*$. By using Theorem 3.12 we can see that the orbit space $\text{InjHom}(K, \text{Mod}_R^*)/\text{Inn}(K)$ of this action is finite. Then we have the following corollary.

Corollary 3.13 *Suppose that $g = 2$ and $p \leq 1$. Let K be a finite index subgroup of Mod_R^* . Then $\text{Out}(K)$ is finite.*

In the other cases, when R has genus at least two and K is a finite index subgroup, we have that $\text{Out}(K)$ is finite as a corollary to the main results in [3], [4]. See [9] for an explicit description of automorphisms of Mod_R^* for a closed surface of genus two.

4 Extending Superinjective Simplicial Maps of $\mathcal{N}(R)$ to Superinjective Simplicial Maps of $\mathcal{C}(R)$

In this section we prove Theorem 1.1 and Theorem 1.2.

Lemma 4.1 *Suppose that $g \geq 2$ and $p \leq 1$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Then λ extends to a superinjective simplicial map $\lambda_* : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$.*

Proof. If x is a nonseparating simple closed curve, we define $\lambda_*([x]) = \lambda([x])$. Let c be a separating simple closed curve on R .

Case 1: Assume that R is closed. Since $g \geq 2$, c separates R into two subsurfaces R_1, R_2 , and both of R_1, R_2 have genus at least one. We take a chain on R_1 , $\{\alpha_1, \dots, \alpha_m\}$ with $i(\alpha_i, \alpha_{i+1}) = 1$, $i(\alpha_i, \alpha_j) = 0$ for $|i - j| > 1$, $[a_i] = \alpha_i \in \mathcal{N}(R)$, as shown in Figure 8, i (for $g = 4$ case when R_1 has genus 3), such that $R_1 \cup \{c\}$ is a regular neighborhood of $a_1 \cup \dots \cup a_n$. Since λ preserves disjointness, nondisjointness and intersection one property, we can see that the chain $\{\alpha_1, \dots, \alpha_n\}$ is mapped by λ into a similar chain, $\{\lambda(\alpha_1), \dots, \lambda(\alpha_m)\}$ with $i(\lambda(\alpha_i), \lambda(\alpha_{i+1})) = 1$, $i(\lambda(\alpha_i), \lambda(\alpha_j)) = 0$ for $|i - j| > 1$. Let $a'_i \in \lambda(\alpha_i)$ such that any two elements in $\{a'_1, \dots, a'_m\}$ have minimal intersection with each other. Let M be a regular neighborhood of $a'_1 \cup \dots \cup a'_n$. Then it is easy to see that M is homeomorphic to $R_1 \cup c$. Let a' be the boundary of M . Suppose that we have another chain on R_1 , $\{\beta_1, \dots, \beta_m\}$ with $i(\beta_i, \beta_{i+1}) = 1$, $i(\beta_i, \beta_j) = 0$ for $|i - j| > 1$, $[b_i] = \beta_i \in \mathcal{N}(R)$ such that $R_1 \cup \{c\}$ is a regular neighborhood of $b_1 \cup \dots \cup b_m$. Again we see that the chain $\{\beta_1, \dots, \beta_m\}$ is mapped by λ into a similar chain, $\{\lambda(\beta_1), \dots, \lambda(\beta_m)\}$ with $i(\lambda(\beta_i), \lambda(\beta_{i+1})) = 1$, $i(\lambda(\beta_i), \lambda(\beta_j)) = 0$ for $|i - j| > 1$. Let $b'_i \in \lambda(\beta_i)$ such that any two elements in $\{b'_1, \dots, b'_m\}$ have minimal intersection with each other. Let T be a regular neighborhood of $b'_1 \cup \dots \cup b'_n$. Then T is homeomorphic to $R_1 \cup c$. Let b' be the boundary of T .

Claim: $[a'] = [b']$.

Proof: We take a similar chain on R_2 , $\{\gamma_1, \dots, \gamma_n\}$ with $i(\gamma_i, \gamma_{i+1}) = 1$, $i(\gamma_i, \gamma_j) = 0$ for $|i - j| > 1$, $[c_i] = \gamma_i \in \mathcal{N}(R)$, such that $R_2 \cup \{c\}$ is a regular neighborhood of $c_1 \cup \dots \cup c_n$. This chain is mapped into a similar chain, $\{\lambda(\gamma_1), \dots, \lambda(\gamma_n)\}$. Let $c'_i \in \lambda(\gamma_i)$ such that any two elements in $\{a'_1, \dots, a'_n, c'_1, \dots, c'_m\}$ and $\{b'_1, \dots, b'_n, c'_1, \dots, c'_m\}$ have minimal intersection with each other. Then a'_i is disjoint from c'_j for any i, j , and b'_i is disjoint from c'_j for any i, j . Then we can choose a regular neighborhood N of $c'_1 \cup \dots \cup c'_n$ in R such that N is disjoint from $M \cup T$. Let c' be the boundary component of N . Then it is easy to see that N is homeomorphic to $R_2 \cup c$ and the boundary components of M and N are isotopic in R . Similarly, the boundary components of T and N are isotopic in R . Hence, $[a'] = [c'] = [b']$. We define $\lambda_*([c]) = [a']$.

Claim: $\lambda_* : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is a simplicial map.

Proof: Let α, β be two vertices in $\mathcal{C}(R)$ such that $i(\alpha, \beta) = 0$. Let x and y be disjoint representatives of α and β respectively. If x, y are two nonseparating simple closed curves then we have $i(\lambda_*(\alpha), \lambda_*(\beta)) = i(\lambda(\alpha), \lambda(\beta)) = 0$. If x is a nonseparating simple closed curve and y is a separating simple closed curve then x lives in a subsurface R_1 which comes from separation by y . Then we could choose a chain as described above which contains x , and then see that by the construction of the image of $[y]$, we get that $i(\lambda_*([x]), \lambda_*([y])) = 0$. If both x and y are separating, then it is easy to see that $R \setminus ((R \setminus x) \cap (R \setminus y))$ has two connected components T_1, T_2 such that T_1 is disjoint from T_2 and y is an essential boundary component of T_1 and x is an essential boundary component of T_2 . Then we see that the




Figure 8: Chains

chains which come from disjoint subsurfaces T_1 and T_2 which will be used to define the images of $[x]$ and $[y]$ are disjoint. Since, λ preserves disjointness we see that the “image” chains will be disjoint, and hence, $i(\lambda_*([x]), \lambda_*([y])) = 0$.

Claim: $\lambda_* : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ is a superinjective simplicial map.

Proof: Let α, β be two vertices in $\mathcal{C}(R)$ such that $i(\alpha, \beta) \neq 0$. Let x and y be representatives of α and β respectively.

(i) Assume that x and y are two nonseparating simple closed curves. Then we have $i(\lambda_*(\alpha), \lambda_*(\beta)) = i(\lambda(\alpha), \lambda(\beta)) \neq 0$ (since λ is superinjective).

(ii) Assume that y is a nonseparating simple closed curve and x is a separating simple closed curve, and x separates R into two subsurfaces R_1, R_2 . W.L.O.G. assume that y is in R_1 . Let $\{\alpha_1, \dots, \alpha_m\}$ be a chain on R_1 with $i(\alpha_i, \alpha_{i+1}) = 1$, $i(\alpha_i, \alpha_j) = 0$ for $|i - j| > 1$, $[a_i] = \alpha_i \in \mathcal{N}(R)$ such that $R_1 \cup \{x\}$ is a regular neighborhood of $a_1 \cup \dots \cup a_m$. Then, since $i(\alpha, \beta) \neq 0$, $i(\beta, \alpha_i) \neq 0$ for some i . Then $i(\lambda_*(\beta), \lambda_*(\alpha_i)) = i(\lambda(\beta), \lambda(\alpha_i)) \neq 0$ (since λ is superinjective). Then it is easy to see that $i(\lambda_*(\alpha), \lambda_*(\beta)) \neq 0$.

(iii) Assume that both x and y are separating and x separates R into two subsurfaces R_1, R_2 . Let $\{\alpha_1, \dots, \alpha_m\}$ be a chain on R_1 with $i(\alpha_i, \alpha_{i+1}) = 1$, $i(\alpha_i, \alpha_j) = 0$ for $|i - j| > 1$, $[a_i] = \alpha_i \in \mathcal{N}(R)$ such that $R_1 \cup \{x\}$ is a regular neighborhood of $a_1 \cup \dots \cup a_m$. Then, since $i(\alpha, \beta) \neq 0$, $i(\beta, \alpha_i) \neq 0$ for some i . Then $i(\lambda_*(\beta), \lambda_*(\alpha_i)) \neq 0$ by (ii). Then it is easy to see that $i(\lambda_*(\alpha), \lambda_*(\beta)) \neq 0$. Hence, we have a superinjective simplicial extension $\lambda_* : \mathcal{C}(R) \rightarrow \mathcal{C}(R)$ of $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$.

Case 2: Assume that R has one boundary component. Then c separates R into two subsurfaces R_1, R_2 . W.L.O.G. assume that R_1 is a genus k subsurface having c as its boundary. We consider chains on R_1 as in the first case, and chains on R_2 such that regular neighborhoods of the curves coming from the chains have c as their essential boundary component and the boundary of R as their inessential boundary component (see Figure 7, ii). Then it is easy to see that the proof of the lemma is similar to case 1. $\square$

Remark: If $g \geq 2$ and $p \geq 2$ and C is the set of separating circles on R which separate R into two pieces such that each piece has genus at least one, then by using chains on these two pieces and following the techniques in the previous lemma we can extend λ to λ_* on C and get a superinjective extension.

Let M be a sphere with k holes and $k \geq 5$. A circle a on M is called an n -circle if a bounds a disk with n holes on M where $n \geq 2$. A *pentagon* in $\mathcal{C}(M)$ is an ordered 5-tuple $(\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5)$, defined up to cyclic permutations, of vertices of $\mathcal{C}(M)$ such that $i(\alpha_j, \alpha_{j+1}) = 0$ for $j = 1, 2, \dots, 5$ and $i(\alpha_j, \alpha_k) \neq 0$ otherwise, where $\alpha_6 = \alpha_1$. A vertex in $\mathcal{C}(M)$ is called an n -vertex if it has a representative which is an n -circle on M .

Let x, y be disjoint simple closed curves on R such that $([x], [y])$ is a peripheral pair, i.e. x, y and a boundary component of R bound a pair of pants on R . $x \cup y$ separate R into two subsurfaces. Let $R_{x,y}$ be the positive genus subsurface of R which comes from this separation. We can identify $\mathcal{N}(R_{x,y})$ with a subcomplex, $L_{x,y}$ of $\mathcal{N}(R)$. By $\lambda_{x,y}$ we will denote the restriction of λ on $\mathcal{N}(R_{x,y})$. If $k : R \rightarrow R$ is a homeomorphism then we will use $k_\#$ for the map induced by k on $\mathcal{N}(R)$ (i.e. $k_\# : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ where $k_\# = [k]$).

An arc i on R is called *properly embedded* if $\partial i \subseteq \partial R$ and i is transversal to ∂R . i is called *nontrivial* (or *essential*) if i cannot be deformed into ∂R in such a way that the end-points of i stay in ∂R during the deformation. If a and b are two disjoint arcs connecting a boundary component of R to itself, a and b are called *linked* if their end points alternate on the boundary component. Otherwise, they are called *unlinked*. The *complex of arcs*, $\mathcal{B}(R)$, on R is an abstract simplicial complex. Its vertices are the isotopy classes of nontrivial properly embedded arcs i in R . A set of vertices forms a simplex if these vertices can be represented by pairwise disjoint arcs. Let i be an essential properly embedded arc on R . Let A be a boundary component of R which has one end point of i and B be the boundary component of R which has the other end point of i . Let N be a regular neighborhood of $i \cup A \cup B$ in R . By Euler characteristic arguments, N is a pair of pants. The boundary components of N are called *encoding circles of i on R* . An essential properly embedded arc i on R is called *type 1* if it joins one boundary component ∂_k of R to itself. i is called *nonseparating* if its complement in R is connected.

The mapping class group, Mod_R , of R is the group of isotopy classes of orientation preserving homeomorphisms of R . The pure mapping class group, $PMod_R$, is the subgroup of Mod_R consisting of isotopy classes of homeomorphisms which preserve each boundary component of R .

Lemma 4.2 *Suppose $g \geq 3$ and $p = 2$. Let $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ be a superinjective simplicial map. Assume that for any peripheral pair $([x], [y])$ on R with x, y disjoint, $\lambda_{x,y}$ agrees with a map, $(g_{x,y})_\# : \mathcal{N}(R_{x,y}) \rightarrow \mathcal{N}(R_{x',y'})$, which is induced by a homeomorphism $g_{x,y} : R_{x,y} \rightarrow R_{x',y'}$ where x', y' are disjoint, $\lambda([x]) = [x']$, $\lambda([y]) = [y']$, $g_{x,y}(x) = x'$, $g_{x,y}(y) = y'$. Then λ agrees with a map $h_\# : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ which is induced by a homeomorphism $h : R \rightarrow R$.*

Proof. Let x, y be disjoint simple closed curves such that $([x], [y])$ is a peripheral pair, and let $(g_{x,y})_\# : \mathcal{N}(R_{x,y}) \rightarrow \mathcal{N}(R_{x',y'})$ be a simplicial map which is induced by a homeomorphism $g_{x,y} : R_{x,y} \rightarrow R_{x',y'}$ where x', y' are disjoint, $\lambda([x]) = [x']$, $\lambda([y]) = [y']$, $g_{x,y}(x) = x'$, $g_{x,y}(y) = y'$ such that $\lambda_{x,y}$ agrees with $(g_{x,y})_\#$ on $\mathcal{N}(R_{x,y})$. Let g be a homeomorphism of R which cuts to a homeomorphism $R_{x,y} \rightarrow R_{x',y'}$ which is isotopic to $g_{x,y}$. Then each homeomorphism of R which cuts to a homeomorphism $R_{x,y} \rightarrow R_{x',y'}$ which is isotopic to $g_{x,y}$, is isotopic to an element in the set $\{gt_x^m t_y^n, m, n \in \mathbb{Z}\}$. It is easy to see that $\lambda_{x,y}$ agrees with the restriction of $(gt_x^m t_y^n)_\#$ on $\mathcal{N}(R_{x,y})$ for all $m, n \in \mathbb{Z}$. Let w be a simple closed curve which is dual to both of x and y (see Figure 9 (i)). Since λ preserves geometric intersection one property by Lemma 2.8, $\lambda([w])$ has a representative which is dual to both of x' and y' . Let P be a regular neighborhood of $x \cup y \cup w$. P is a genus one surface with two boundary components. Let t be the essential boundary component of P . Let Q be the genus one subsurface with two boundary components of R which has $g(t)$ as its boundary. Then by




Figure 9: (i) A dual curve, (ii) Pants decomposition

using the properties of λ it is easy to see that $\lambda([w])$ has a representative which lies in the interior of Q , and dual to both of x' and y' and there exists $m_o, n_o \in \mathbb{Z}$ such that $gt_x^{m_o} t_y^{n_o}$ agrees with λ on $[w]$. Let $D_{x,y}$ be the set of isotopy classes of simple closed curves which are dual to each of x and y on R .

Claim 1: $(gt_x^{m_o} t_y^{n_o})_{\#}$ agrees with λ on $\{[x]\} \cup \{[y]\} \cup L_{x,y} \cup D_{x,y}$.

Proof: It is clear that $(gt_x^{m_o} t_y^{n_o})_{\#}([x]) = \lambda([x]) = [x']$ and $(gt_x^{m_o} t_y^{n_o})_{\#}([y]) = \lambda([y]) = [y']$. Since $\lambda_{x,y}$ agrees with the restriction of $(gt_x^{m_o} t_y^{n_o})_{\#}$ on $\mathcal{N}(R_{x,y})$, $(gt_x^{m_o} t_y^{n_o})_{\#}$ agrees with λ on $L_{x,y}$. We have seen that $(gt_x^{m_o} t_y^{n_o})_{\#}$ agrees with λ on $[w]$. Let w_1 be a simple closed curve which is disjoint from w and dual to both of x and y . As we described before, there exists $\tilde{m}, \tilde{n} \in \mathbb{Z}$ such that λ agrees with $(gt_x^{\tilde{m}} t_y^{\tilde{n}})_{\#}$ on $[w_1]$. Since w and w_1 are disjoint nonseparating curves, $i(\lambda([w]), \lambda([w_1])) = 0$. Then by using the properties that λ preserves disjointness and nondisjointness, we can see that $m = \tilde{m}$ and $n = \tilde{n}$. This shows that $(gt_x^{m_o} t_y^{n_o})_{\#}$ also agrees with λ on $[w_1]$. Given any simple closed curve v which is dual to both of x and y , we can find a sequence of dual curves to both of x and y , connecting w to v , such that each consecutive pair is disjoint, i.e. the isotopy classes of these curves define a path between w and v in $\mathcal{N}(R)$. Then using the argument given above and the sequence, we conclude that $(gt_x^{m_o} t_y^{n_o})_{\#}$ agrees with λ on $D_{x,y}$. Hence, $(gt_x^{m_o} t_y^{n_o})_{\#}$ agrees with λ on $\{[x]\} \cup \{[y]\} \cup L_{x,y} \cup D_{x,y}$. This proves claim 1. Let $h_{x,y} = gt_x^{m_o} t_y^{n_o}$. We have that $(h_{x,y})_{\#}$ agrees with λ on $\{[x]\} \cup \{[y]\} \cup L_{x,y} \cup D_{x,y}$.

We complete x, y to a pair of pants decomposition P consisting of nonseparating circles as shown in Figure 9 (ii) for $g = 3, p = 2$, (similar configurations can be chosen for $g \geq 3$). Let t, z, w, c, k, r be as shown in the figure.

Claim 2: $(h_{x,y})_{\#}$ agrees with λ on c .

Proof: Let t', z', x', y', k', r' be pairwise disjoint representatives of $\lambda([t]), \lambda([z]), \lambda([x]), \lambda([y]), \lambda([k]), \lambda([r])$ respectively and let w' be a representative of $\lambda([w])$ which has minimal intersection with each of t', z', x', y', k', r' . Since k, r, z, y, ∂_1 bound a sphere with five holes, T , on R , containing x and c , then k', r', z', y' and a boundary component of R bound a sphere with five holes, T' , on R by Lemma 2.6. Since x and c have geometric intersection 2 and algebraic intersection 0 in T , it is easy to see that there exist vertices $\gamma_1, \gamma_2, \gamma_3$ of $\mathcal{C}(T)$ such that $(\gamma_1, \gamma_2, [x], \gamma_3, [c])$ is a pentagon in $\mathcal{C}(T)$, γ_1 and γ_3 are 2-vertices, γ_2 is a 3-vertex, and $\{[x], \gamma_3\}$, $\{[x], \gamma_2\}$, $\{[c], \gamma_3\}$ and $\{\gamma_1, \gamma_2\}$ are codimension-zero simplices of $\mathcal{C}(T)$, and each of γ_i has a representative which is nonseparating on R . Since λ is superinjective, we




Figure 10: Arcs and their encoding circles

can see that $(\lambda(\gamma_1), \lambda(\gamma_2), \lambda([x]), \lambda(\gamma_3), \lambda([c]))$ is a pentagon in $\mathcal{C}(T')$. By Lemma 2.6, $\lambda(\gamma_1)$ and $\lambda(\gamma_3)$ are 2-vertices, and $\lambda(\gamma_2)$ is a 3-vertex in $\mathcal{C}(T')$. Since λ is an injective simplicial map $\{\lambda([x]), \lambda(\gamma_3)\}$, $\{\lambda([x]), \lambda(\gamma_2)\}$, $\{\lambda([c]), \lambda(\gamma_3)\}$ and $\{\lambda(\gamma_1), \lambda(\gamma_2)\}$ are codimension-zero simplices of $\mathcal{C}(T')$. It is easy to see that $\lambda([c])$ has a representative which is disjoint from t', z', y' . Then $\{\lambda([x]), \lambda([c])\}$ have representatives with geometric intersection 2 and algebraic intersection 0 in the sphere with four holes bounded by t', z', y' and the boundary component of R , [8]. We also have that $(h_{x,y})_{\#}([c])$ has a representative c'' which is disjoint from t', z', y', w' and which intersects x' geometrically twice and algebraically 0 times. Since $(h_{x,y})_{\#}$ agrees with λ on $\{[x]\} \cup \{[y]\} \cup L_{x,y} \cup D_{x,y}$ it is easy to see that $[c'] = [c'']$, i.e. $(h_{x,y})_{\#}$ agrees with λ on c .

Claim 3: $(h_{x,y})_{\#}$ agrees with λ on the class of every nonseparating circle on R .

Proof: Let z be a nonseparating simple closed curve on R . Let t be another simple closed curve disjoint from z such that z, t and ∂_1 bound a pair of pants in R . Let i and j be nonseparating type 1 arcs connecting ∂_1 to itself such that i has x, y as its encoding circles and j has z, t as its encoding circles. W.L.O.G. we can assume that i and j have minimal intersection. By Lemma 3.8 in [4], there is a sequence $i = r_0 \rightarrow r_1 \rightarrow \dots \rightarrow r_{n+1} = j$ of essential properly embedded nonseparating type 1 arcs joining ∂_1 to itself so that each consecutive pair is disjoint, i.e. the isotopy classes of these arcs define a path in $\mathcal{B}(R)$ between i and j . Let x_i, y_i be the encoding circles for r_i for $i = 1, \dots, n$. For the pair of arcs i and r_1 , we will consider the following cases:

Case i: Assume that i and r_1 are linked, i.e. their end points alternate on the boundary component ∂_1 . Then a regular neighborhood of $i \cup r_1 \cup \partial_1$ is a genus one surface with two boundary components N , and the arcs i, r_1 and their encoding circles x, y, x_1, y_1 on N are as shown in Figure 10 (i). In this case, we complete $\{x, y, x_1, y_1\}$ to a curve configuration G consisting of nonseparating circles which is shown in Figure 11 (i), for $g = 3, p = 3$, see [7], such that the isotopy classes of Dehn twists about the elements of this set generate $PMod_R$ and all the curves in this set are (i) either disjoint from x, y or simultaneous dual to x, y , and (ii) either disjoint from x_1, y_1 or simultaneous dual to x_1, y_1 . Then, since all the curves in G are either disjoint from x, y or simultaneous dual to x, y , by claim 1 we have that $(h_{x,y})_{\#}([x]) = \lambda([x])$ for every $x \in G$. Similarly, since all the curves in G are either disjoint from x_1, y_1 or simultaneous dual to x_1, y_1 , by claim 1 we have $(h_{x_1,y_1})_{\#}([x]) = \lambda([x])$ for every $x \in G$. Hence $(h_{x,y})_{\#}([x]) = \lambda([x]) = (h_{x_1,y_1})_{\#}([x])$ for every $x \in G$. Then $(h_{x,y}^{-1} h_{x_1,y_1})_{\#} \in C_{Mod_R}(PMod_R)$. By Theorem 5.3 in [7], $C_{Mod_R}(PMod_R) = \{1\}$. Hence $(h_{x,y})_{\#} = (h_{x_1,y_1})_{\#}$.




Figure 11: Encoding circles in a curve configuration

Case ii: Assume that i and r_1 are unlinked, i.e their end points don't alternate on the boundary component ∂_1 . Then a regular neighborhood of $i \cup r_1 \cup \partial_1$ is a sphere with four boundary components S_4^2 , and the arcs i, r_1 and their encoding circles x, y, x_1, y_1 on S_4^2 are as shown in Figure 10 (ii). Let w be the boundary component of S_4^2 which is different from x_1, y, ∂_1 . If w is a nonseparating curve, then we complete $\{x, y, x_1, y_1\}$ to a curve configuration G consisting of nonseparating circles as shown in Figure 11 (ii) such that the isotopy classes of Dehn twists about the elements of G generate $PMod_R$. By claim 1 and claim 2, $(h_{x,y})_\#([x]) = \lambda([x])$ for every $x \in G$ and $(h_{x_1,y_1})_\#([x]) = \lambda([x])$ for every $x \in G$. Then $(h_{x,y}^{-1}h_{x_1,y_1})_\#([x]) = [x]$ for every $x \in G$. Then $(h_{x,y}^{-1}h_{x_1,y_1})_\# \in C_{Mod_R}(PMod_R)$. By Theorem 5.3 in [7], $C_{Mod_R}(PMod_R) = \{1\}$. Hence $(h_{x,y})_\# = (h_{x_1,y_1})_\#$.

Suppose that w is a separating curve. By the remark after Lemma 4.1, we can extend λ to a superinjective map λ_* on a subcomplex of $\mathcal{C}(R)$ containing separating circles on R which separate R into two pieces such that each piece has genus at least one. Notice that w is such a circle. For the rest of the proof, we will use λ for this extension. We will do the case when x_1 and y are not isotopic. The other case can be done similarly. Let M be the subsurface which has w on its boundary and which does not contain N . Let T be the closure of $R \setminus \{M \cup N\}$. The circles x_1 and y are boundary components of T . Since w is an essential separating circle and $p = 2$, M has genus at least one. In Figure 12, we show M, N, T for a special case. By claim 1, $(h_{x,y})_\#([x]) = \lambda([x])$ for every $x \in \mathcal{N}(T)$. Similarly, $(h_{x_1,y_1})_\#([x]) = \lambda([x])$ for every $x \in \mathcal{N}(T)$. Then $(h_{x,y}^{-1}h_{x_1,y_1})_\#([x]) = [x]$ for every $x \in \mathcal{N}(T)$. Then the restriction of $(h_{x,y}^{-1}h_{x_1,y_1})_\#$ on $\mathcal{N}(T)$ is in $C(PMod_T)$. By Theorem 5.3 in [7], $C(PMod_T) = \{1\}$. Hence $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\mathcal{N}(T)$. It is easy to see that $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on the set $\{[x_1], [w], [y]\}$. Following the proof of claim 2 (considering that we have the extended superinjective simplicial map on “good” separating circles), we see that $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\{[x], [y_1]\}$. Then, since Dehn twists about x and y_1 generate $PMod_N$, the restriction of $(h_{x,y}^{-1}h_{x_1,y_1})_\#$ on $\mathcal{C}(N)$ is in $C(PMod_N)$. By Theorem 5.3 in [7], $C(PMod_N) = \{1\}$. Hence $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\mathcal{C}(N)$. By claim 1, $(h_{x,y}^{-1}h_{x_1,y_1})_\#([x]) = [x]$ for every $x \in \mathcal{N}(M)$. Then the restriction of $(h_{x,y}^{-1}h_{x_1,y_1})_\#$ on $\mathcal{N}(M)$ is in $C(PMod_M)$. By considering the action on oriented circles and using Theorem 5.3 in [7], we see that $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\mathcal{N}(M)$. So we have $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\mathcal{N}(M) \cup \mathcal{C}(N) \cup \mathcal{N}(T)$. In Figure 12 we see the curve c which is dual to each of x, y, x_1, y_1 . By claim 1, $(h_{x,y})_\#([c]) = \lambda([c]) = (h_{x_1,y_1})_\#([c])$. Then, since $(h_{x,y})_\#$ and $(h_{x_1,y_1})_\#$ agree on $[c]$, we have $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ on $\mathcal{N}(R)$. Then, since $g \geq 3$, $(h_{x,y})_\# = (h_{x_1,y_1})_\#$. If w is a boundary component of R , then showing that $(h_{x,y})_\# = (h_{x_1,y_1})_\#$ is similar.




Figure 12: Curves intersecting twice

In both cases we have seen that $(h_{x,y})_{\#} = (h_{x_1,y_1})_{\#}$. By using our sequence, with an inductive argument we have $(h_{x,y})_{\#} = (h_{z,t})_{\#}$, and $(h_{x,y})_{\#} = (h_{z,t})_{\#} = \lambda$ on $\{[x]\} \cup \{[y]\} \cup \{[z]\} \cup \{[t]\} \cup L_{x,y} \cup D_{x,y} \cup L_{z,t} \cup D_{z,t}$. In particular we see that $(h_{x,y})_{\#}$ agrees with λ on any nonseparating curve z , hence $(h_{x,y})_{\#}$ agrees with λ on $\mathcal{N}(R)$. This proves the lemma. $\square$

Theorem 4.3 *Suppose that the genus of R is at least two and R has at most $g-1$ boundary components. A simplicial map $\lambda : \mathcal{N}(R) \rightarrow \mathcal{N}(R)$ is superinjective if and only if λ is induced by a homeomorphism of R .*

Proof. If λ is induced by a homeomorphism of R , then it preserves disjointness and nondisjointness and hence it is superinjective. Assume that λ is superinjective. In the cases when R is a closed surface or when R has exactly one boundary component, by Lemma 4.1 λ extends to a superinjective simplicial map λ_* on $\mathcal{C}(R)$. Then λ_* is induced by a homeomorphism $h : R \rightarrow R$, i.e. $\lambda_*(\alpha) = h_{\#}(\alpha)$ for each vertex α in $\mathcal{C}(R)$, by the main results in [3] and [4] and by Theorem 1.3. Hence λ is induced by the homeomorphism h .

Assume that $g \geq 3$ and $p = 2$. Let x and y be disjoint nonseparating circles such that x, y and a boundary component, ∂_1 , of R bound a pair of pants on R . Let x', y' be disjoint representatives of $\lambda([x]), \lambda([y])$ respectively. By using Lemma 2.5 and knowing that λ preserves disjointness and nondisjointness, it is easy to see that λ maps $\mathcal{N}(R_{x \cup y})$ to $\mathcal{N}(R_{x' \cup y'})$. Since every essential separating curve on $R_{x \cup y}$ separates R into two subsurfaces each of which has genus at least one, using chains on these subsurfaces we could extend λ to a superinjective simplicial map $\lambda_{x,y} : \mathcal{C}(R_{x \cup y}) \rightarrow \mathcal{C}(R_{x' \cup y'})$ as in case 1 in Lemma 4.1. Then by the main results in [4], there exists a homeomorphism $h : R_{x \cup y} \rightarrow R_{x' \cup y'}$ such that $h(x) = x', h(y) = y'$ and $\lambda_{x,y}$ is induced by h . Then the proof of the theorem follows from Lemma 4.2.

Now assume that $g \geq 4$ and $3 \leq p \leq g-1$. We will give the proof when $p = g-1$. The proof of the remaining cases is similar. Let $\{a_1, \dots, a_{2(p-1)}\}$ be a set of pairwise disjoint nonseparating circles such that (a_{2i+1}, a_{2i+2}) is a peripheral pair as shown in Figure 13 for $i = 0, \dots, p-2$. Let $R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ be the genus two surface with $2p-1$ boundary components which comes from the separation by $a_1 \cup a_2 \dots \cup a_{2(p-1)}$. Exactly one of the boundary components of $R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ is a boundary component of R . We identify $\mathcal{N}(R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}})$ with a subcomplex $L_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ of $\mathcal{N}(R)$. Let $\lambda_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ denote the restriction of λ on $\mathcal{N}(R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}})$. Let $a'_1, \dots, a'_{2(p-1)}$ be pairwise disjoint representatives of $\lambda([a_1]), \dots, \lambda([a_{2(p-1)}])$ respectively. By using Lemma 2.5 and the properties of λ , it is easy to see that λ maps $\mathcal{N}(R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}})$ to $\mathcal{N}(R_{a'_1 \cup a'_2 \dots \cup a'_{2(p-1)}})$. Since every essential separating curve on $R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ separates R into two subsurfaces each of




Figure 13: Cutting R along peripheral pairs

which has genus at least one, using chains on these subsurfaces we extend λ to a superinjective simplicial map $\lambda_{a_1 \cup a_2 \dots \cup a_{2(p-1)}} : \mathcal{C}(R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}) \rightarrow \mathcal{C}(R_{a'_1 \cup a'_2 \dots \cup a'_{2(p-1)}})$ as in Lemma 4.1. Then by the main results in [4], there exists a homeomorphism $h : R_{a_1 \cup a_2 \dots \cup a_{2(p-1)}} \rightarrow R_{a'_1 \cup a'_2 \dots \cup a'_{2(p-1)}}$ such that $h(a_i) = a'_i$ and $\lambda_{a_1 \cup a_2 \dots \cup a_{2(p-1)}}$ is induced by h . If we replace (a_{2p-3}, a_{2p-2}) with another peripheral pair (b_{2p-3}, b_{2p-2}) where each of b_{2p-3} and b_{2p-2} is disjoint from $a_1 \cup a_2 \dots \cup a_{2p-4}$, then by similar techniques we get a homeomorphism $t : R_{a_1 \cup a_2 \dots \cup a_{2p-4} \cup b_{2p-3} \cup b_{2p-2}} \rightarrow R_{a'_1 \cup a'_2 \dots \cup a'_{2p-4} \cup b'_{2p-3} \cup b'_{2p-2}}$ such that $t(a_i) = a'_i$, $t(b_j) = b'_j$, where b'_{2p-3}, b'_{2p-2} are pairwise disjoint representatives of $\lambda([b_{2p-3}]), \lambda([b_{2p-2}])$ respectively, and the map $\lambda_{a_1 \cup a_2 \dots \cup a_{2p-4} \cup b_{2p-3} \cup b_{2p-2}}$ is induced by t . Then by following the techniques in the proof of Lemma 4.2, we see that there exists a homeomorphism $r : R_{a_1 \cup a_2 \dots \cup a_{2p-4}} \rightarrow R_{a'_1 \cup a'_2 \dots \cup a'_{2p-4}}$ such that $r(a_i) = a'_i$ and $\lambda_{a_1 \cup a_2 \dots \cup a_{2p-4}}$ is induced by r . Then by an inductive argument, there exists a homeomorphism $q : R \rightarrow R$ such that λ is induced by q . $\square$

Now we consider the graph $\mathcal{G}(R)$ defined by Schaller. The vertex set of $\mathcal{G}(R)$ is the set of isotopy classes of nontrivial nonseparating simple closed curves on R . Two vertices are connected by an edge if and only if their geometric intersection number is one. In the proof of the following theorem, we use some of Schaller's results given in [13] that if $g \geq 2$ and R is not a closed surface of genus two, then $\text{Aut}(\mathcal{G}(R)) = \text{Mod}_R^*$.

Theorem 4.4 *Suppose that R has genus at least two. If R is not a closed surface of genus two, then $\text{Aut}(\mathcal{N}(R)) = \text{Mod}_R^*$. If R is a closed surface of genus two, then $\text{Aut}(\mathcal{N}(R)) = \text{Mod}_R^* / \mathcal{C}(\text{Mod}_R^*)$.*

Proof. Assume that R is not a closed surface of genus two. If $[f] \in \text{Mod}_R^*$, then $[f]$ induces an automorphism of $\mathcal{N}(R)$. If $[f]$ fixes the isotopy class of every nontrivial nonseparating simple closed curve on R , then f is orientation preserving and it can be shown that it is isotopic to id_R . An automorphism λ of $\mathcal{N}(R)$ is a superinjective simplicial map of $\mathcal{N}(R)$. By Lemma 2.8, it preserves geometric intersection one property, and hence induces an automorphism of $\mathcal{G}(R)$. An automorphism of $\mathcal{G}(R)$ is induced by a homeomorphism of R , [13] (Note that if R has at most $g - 1$ boundary components, then Theorem 4.3 also implies that λ is induced by a homeomorphism of R). Then it is easy to see that we get $\text{Aut}(\mathcal{N}(R)) = \text{Mod}_R^*$. Assume that R is a closed surface of genus two. If $[f] \in \text{Mod}_R^*$, then $[f]$ induces an automorphism of $\mathcal{N}(R)$. If $[f]$ fixes the isotopy class of every nonseparating simple closed curve on R then f is either isotopic to id_R or the hyperelliptic involution. An automorphism of $\mathcal{N}(R)$ is a superinjective simplicial map of $\mathcal{N}(R)$ and by Theorem 4.3 it is induced by a homeomorphism of R . Then we have $\text{Aut}(\mathcal{N}(R)) = \text{Mod}_R^* / \mathcal{C}(\text{Mod}_R^*)$. $\square$

Note that by the main results of this paper and the results in [3], [4], [5], [13], we have that if R has genus at least two, then $\text{Aut}(\mathcal{N}(R)) = \text{Aut}(\mathcal{C}(R)) = \text{Aut}(\mathcal{G}(R))$.

Acknowledgments

We thank John D. McCarthy for his suggestions and many discussions about this work. We also thank Peter Scott for his comments.

References

- [1] J.S. Birman, A. Lubotzky, J.D. McCarthy, Abelian and solvable subgroups of the mapping class groups, *Duke Math. Jour.* 50 (4) (1983), 1107-1120.
- [2] W.J. Harvey, Geometric structures of surface mapping class groups, *Homological Group Theory* (C.T. Wall, ed), *London Math. Soc. Lecture Notes*, No.36, Cambridge Univ. Press, London, 1979, 255-269.
- [3] E. Irmak, Superinjective simplicial maps of complexes of curves and injective homomorphisms of subgroups of mapping class groups, *Topology* 43 (2004) 513-541.
- [4] E. Irmak, Superinjective simplicial maps of complexes of curves and injective homomorphisms of subgroups of mapping class groups II, submitted, available at <http://front.math.ucdavis.edu/math.GT/0311407>.
- [5] N.V. Ivanov, Automorphisms of complexes of curves and of Teichmuller spaces, *International Mathematics Research Notices*, No.14 (1997), 651-666.
- [6] N.V. Ivanov, Subgroups of Teichmuller modular groups, *Translations of Mathematical Monographs*, 115, AMS, 1992.
- [7] N.V. Ivanov, J.D. McCarthy, On Injective Homomorphisms between Teichmuller Modular Groups I, *Inventiones Mathematicae* 135, 425-486 (1999).
- [8] M. Korkmaz, Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology and its Applications*, 95 (2) (1999), 85-111.
- [9] J.D. McCarthy, Automorphisms of surface mapping class groups. A recent theorem of N.Ivanov, *Inventiones Mathematicae* 84, 49-71 (1986).
- [10] J.D. McCarthy, Normalizers and centralizers of pseudo-Anosov mapping classes. The manuscript is available for informal distribution, on request.
- [11] F. Luo, On Non-separating simple closed curves in a compact surface, *Elect. Res. Announc. of AMS*, V.1, (1995) 1.
- [12] L. Mosher, Tiling the projective foliation space of a punctured surface, *Transactions of the American Mathematical Society*, 306 (1) (1988), 1-70.
- [13] P.S. Schaller, Mapping class groups of hyperbolic surfaces and automorphism groups of graphs, *Composito Mathematica* 122 (2000), 243-260.

University of Michigan, Department of Mathematics, Ann Arbor, MI 48109, USA;
eirmak@umich.edu